

Further details to

Schänis, Switzerland: Long-term forest meteorological data from the Long-term Forest Ecosystem Research Programme (LWF), from 1997-2016

Matthias Haeni^{1,*}, Arthur Gessler¹, Martine Rebetez¹

¹ Swiss Federal Institute for Forest, Snow and Landscape Research WSL

November 2016

Summary

High quality meteorological data are needed for long-term forest ecosystem research, particularly in the light of global change. The long-term data series published here comprises almost 20 years of two meteorological stations in Schänis in Switzerland where one station is located within a natural mixed forest stand (SCB) with European beech (*Fagus sylvatica*; 130-150 yrs), European silver fir (*Abies alba*; 130-150 yrs) and European ash (*Fraxinus excelsior*; 130-150 yrs) as dominant tree species. A second station is situated in the very vicinity outside of the forest (field station, SCF). The meteorological time series are presented in hourly time resolution of air temperature, relative humidity, precipitation, photosynthetically active radiation (PAR) and wind speed. Schänis is part of the Long-term Forest Ecosystem Research Programme (LWF) established and maintained by the Swiss Federal Research Institute WSL.

*Correspondence to Matthias Haeni, matthias.haeni@wsl.ch

1 Introduction

High quality meteorological data often found the basis for numerous scientific analyses in the light of global change. The Long-term Forest Ecosystem Research programme (LWF) at the Swiss Federal Research Institute WSL investigates anthropogenic and natural forces on the long-term health of the forests and the potential risks to humans. Furthermore, a profound understanding of the relationship between processes, causes and effects in the forest ecosystem are necessary (Schaub et al.; 2011) and weather data usually found the basis of the input factors of such processes.

Within the LWF, measurements have been started in 1995 on 16 forest stations within the level 2 plots of ICP Forests. This data repository here compromises the most important meteorological time series of one of the stations: Schänis, located in the canton of St. Gallen in Switzerland.

2 Materials and Methods

2.1 Site

The measurements were carried out at Schänis in the Swiss Canton of St. Gallen, located between the Lake of Zürich and the Walensee (Fig. 1). Time series of two plots have been recorded, one within the forest (Fig. 2a) and a second one outside of the forest (field station, Fig. 2b). Tab. 1 shows further details on the research station characteristics.




Figure 1: Site aerial photo from both Schänis stations. Source: Google. 2016. Site aerial photo from both Schänis stations. Retrieved November 11, 2016, from <https://www.google.ch/maps>

Table 1: Meteorological station setting

Community / canton	Schänis / SG	
Commissioning date	17 September 1997	
	SCB (forest)	SCF (field)
Coordinates (Lat. / Lon.)	47.165937 ° / 9.066649 °	47.151248 ° / 9.059543 °
Altitude	698 m a.s.l.	627 m a.s.l.
Orientation	W	W
Mean slope	48.8 %	45.8 %

**(a)** SCB: Forest station**(b)** SCF: Field station**Figure 2:** Images of the meteo stations in Schänis.**Table 2:** Tree stand characteristics

Woodland association*	Nr. 13; Cardamino-Fagetum tilietosum
Main tree species (lat.)	<i>Fagus sylvatica</i>
Management system	High forest
Silvicultural system	Group selection
Top-height diameter	55.8 cm
Plot size	2 ha
Number of trees DBH ≥ 12 cm (2011)	N = 611
Maximum tree age	
<i>Fagus sylvatica</i>	130-150 yrs
<i>Abies alba</i>	130-150 yrs
<i>Fraxinus excelsior</i>	130-150 yrs
Geology (German)	Tertiär. Subalpine Molasse
	Oligocaen, Chattien, Kalknagelfluh
Soil types (WSL)	Pseudovergleyte Braunerde

*after [Ellenberg and Klötzli \(1972\)](#)

2.2 Sensors

Sensor and measurement specifications

Each station within the LWF acquires data continuously over ten minute periods. Here with this data set, hourly data is provided. All measurements are post-measurement values (e.g. values from 15:10 o'clock compromise data averaged over the period of 15:00:01 to 15:10:00). The sensor types are listed in Tab. 3.

Table 3: Sensor specifications

Parameter	Unit	Height	Interval	Mode	Producer / Model	Fig.
Temperature	°C	2 m	10 min	Mean	Retronic / MP100A	3
Rel. humidity	%	2 m	10 min	Mean	Retronic / MP100A	3
Precipitation	$mm\ m^{-2}$	1.5 m	10 min	Sum	Campbell Sc. / ARG100	4
PAR*	$W\ m^{-2}$	3 m	10 min	Mean	Skye Instr. / Quantum	5
Wind speed	$m\ s^{-1}$	4.6 m	10 min	Mean	Vector Instr. / A100R	6

Annotations

Temperature / relative humidity	For both temperature and relative humidity a combined sensor is used. The thermometer is a commonly used DIN Class B sensor (accuracy ± 0.3 K) whereas the hygrometer has an accuracy of ± 1.5 % rH (both at $23 \pm 5^\circ\text{C}$).
Precipitation	Precipitation is collected by a gauge with a funnel width of 254 mm and a rocker switch registering 2 mm of precipitation.
PAR	The PAR sensor measures solar radiation with wavelength between 400 nm and 700 nm.
Wind speed	Wind speed is measured by an anemometer with a resolution of 1.25 m.

*Photosynthetically active radiation

3 Results

3.1 Air temperature (temp)

The recorded data from 10 minute intervals have been averaged to the full hour (minutes 30, 40, 50, 00, 10, 20 averaged). First, minor manual data exclusions have been made where data values were in irrecoverably bad shape. Then an absolute filter ($-42^{\circ}\text{C} < x < +47^{\circ}\text{C}$) and a relative filter have been applied (0.001 % quantile). A differential filter removed data entries with larger jumps than 6°C from one hour to the other. Then a relative moving window filter has been applied, where data points have been excluded if they exceeded the 0.1 % quantiles on both ends (0.1 %, > 99.9 %) in a fourteen day moving window for every single hour. After filling small gaps linearly (two hours), we compared the data from the individual station with the partner station (field and forest stations). Large differences between the two time series ($> 20^{\circ}\text{C}$) were removed. Furthermore, in a density plot between field and forest station, data outside of the 99.9 % contour (not including the extreme edges) were removed (for details please contact Matthias Haeni, matthias.haeni@wsl.ch). If the station data correlated in a linear manner with an $R^2 > 0.95$, data gaps were filled with the calculated data from the partner station.

3.2 Relative humidity (rH)

The recorded data from 10 minute intervals have been averaged to the full hour (minutes 30, 40, 50, 00, 10, 20 averaged). First, minor manual data exclusions have been made where data values were in irrecoverably bad shape. Then an absolute filter ($0 \% < x < 110 \%$) and a relative filter have been applied (0.001 % quantile). A differential filter removed data entries with larger jumps than 30 % from one hour to the other. The data have then been compared to Meteonorm computed data (Meteonorm; 2016; <http://www.meteonorm.com/en/>), and the time series have been scaled conservatively for each individual year to counteract large sensor drifts. Then, a relative moving window filter has been applied, where data points have been excluded if they exceeded the 0.1 % quantiles on both ends ($< 0.1 \%$, $> 99.9 \%$) in a fourteen day moving window for every single hour. After filling small gaps linearly (two hours), we compared the data from the individual station with the partner stations (field to forest stations and vice versa). Large differences between the two time series ($> 60 \%$) were removed. Furthermore, in a density plot between field and forest station, data outside of the 99.9 % contour not including the extreme edges were removed (for details please contact Matthias Haeni, matthias.haeni@wsl.ch). If the station data correlated in a linear manner with an $R^2 > 0.95$, data gaps were filled with the calculated data from the partner station.

3.3 Precipitation (precip)

The recorded data from 10 minute intervals have been summed up to the full hour (minutes 10, 20, 30, 40, 50, 00 to the hour). First, minor manual data exclusions have been made where data values were in irrecoverably bad shape. Then an absolute filter ($0 \text{ mm} < x < 100 \text{ mm}$) has been applied. Long periods of not precipitation (> 120 days of zero values) have been excluded, because in this case a clogged rain gauge system must be assumed. Then we compared the data from the individual station with the partner stations (field to forest stations and vice versa). Large differences between the two time series ($> 50 \text{ mm}$) were removed. Furthermore, in a density plot between field and forest station, data outside of the 99.9 % contour not including the extreme edges were removed (for details please

contact Matthias Haeni, matthias.haeni@wsl.ch). If the station data correlated in a linear manner with an $R^2 > 0.8$, data gaps were filled with the calculated data from the partner station.

3.4 Photosynthetically active radiation (PAR)


The recorded data from 10 minute intervals have been averaged to the full hour (minutes 30, 40, 50, 00, 10, 20 averaged). First, minor manual data exclusions have been made where data values were in irrecoverably bad shape. Then an absolute filter ($0 \text{ W m}^{-2} < x < 2500 \text{ W m}^{-2}$) has been applied. After filling small gaps linearly (two hours), we conducted a so-called night shift correction. Night time hours have been calculated in two ways: 1) From the data, where in the morning average PAR was first above 100 W m^{-2} and in the evening first below 100 W m^{-2} . To the first so-called 'sunrise' time, half an hour was subtracted and to the latter so-called 'sunset' time, half an hour was added. 2) Sunset and sunrise times have been calculated from the google maps API (Google; 2016; <https://developers.google.com/maps/>). From both the calculated methods, the more conservative 'sunrise' and 'sunset' times were chosen, i.e. the earlier one for 'sunrise' and the later one for 'sunset'. For the hours in between the 'sunset' and 'sunrise' times, for these night time hours, no sunlight conditions should lead to zero W m^{-2} PAR. The average deviations of these night time hours were then corrected for each individual day. Data were then scaled from year to year by comparing to the average 99 % density contour of all years and scaling conservatively to counteract large sensor drift (for details please contact Matthias Haeni, matthias.haeni@wsl.ch).

3.5 Wind speed (ws)


The recorded data from 10 minute intervals have been averaged to the full hour (minutes 30, 40, 50, 00, 10, 20 averaged). First, minor manual data exclusions have been made where data values were in irrecoverably bad shape. Then an absolute filter ($0 \text{ m s}^{-1} < x < 100 \text{ m s}^{-1}$) has been applied. Then, we compared the data from the individual station with the partner station (field to forest stations and vice versa). In a density plot between field and forest station, data outside of the 99.9 % contour not included the extreme edges were removed (for details please contact Matthias Haeni, matthias.haeni@wsl.ch). If the station data correlated in a linear manner with an $R^2 > 0.8$, data gaps were filled with the calculated data from the partner station.

The data are shown on the next two pages.

SCB



SCF



References

Ellenberg, H. and Klötzli, F. (1972). Die waldgesellschaften der schweiz, *Mitt. Schweiz. Anst. forstl. Versuchsw* **48**(4): 930.

Google (2016). Google Maps API.

URL: <https://developers.google.com/maps/>

Meteonorm (2016). Irradiation data for every place on Earth, easily accessible with Meteonorm.

URL: <http://www.meteonorm.com/en/>

Schaub, M., Dobbertin, M., Kräuchi, N. and Dobbertin, M. K. (2011). Preface: long-term ecosystem research: understanding the present to shape the future, *Environmental monitoring and assessment* **174**(1): 1–2.

Appendix

Pictures of sensors



Figure 3: Combined air temperature and humidity sensor



Figure 4: Precipitation sensor



Figure 5: PAR sensor



Figure 6: Wind speed sensor